CLAIMS:

5


10

15

20

25

1. A compound of the formula:



wherein R^1 is independently in each occurrence i) a C_{1-40} hydrocarbyl group, ii) a C_{1-40} hydrocarbyl group wherein one or more carbons are substituted by one or more heteroatoms selected from S, N, O, P, B or Si atoms, or iii) a halogenated derivative of iii) or iv), with the proviso that in at least one occurrence, R^1 is crosslinkable group;

 R^2 is independently in each occurrence hydrogen, halogen, C_{1-20} hydrocarbyl, C_{1-20} hydrocarbyloxy, C_{1-20} thioether, C_{1-20} hydrocarbylcarbonyloxy, di(C_{1-20} hydrocarbyl)amino, or cyano;

 Ar^1 , Ar^2 , Ar^3 and Ar^4 are independently in each occurrence C_{6-20} aromatic groups, optionally containing one or more S, N, O, P, B or Si heteroatoms, or a halo-, C_{1-20} hydrocarbyl-, $di(C_{1-20}$ hydrocarbyl)amino-, C_{1-20} hydrocarbyloxy-, $tri(C_{1-10}$ hydrocarbyl)silvy-, or $tri(C_{1-10}$ hydrocarbyl)silvy- substituted derivative thereof;

a and b independently in each occurrence are 0 or 1; and

 X^1 and X^2 independently in each occurrence are a covalent bond, O, S, SO₂, CH₂, C(R³)₂ or NR³, wherein R³ is selected from the group consisting of C₁₋₂₂ alkyl, C₁₋₂₂ cycloalkyl, C₆₋₂₄ aryl, and C₇₋₂₄ aralkyl.

- 2. A compound according to claim 1 wherein R^1 independently each occurrence is selected from the group consisting of C_{1-40} hydrocarbyl, C_{3-40} hydrocarbyl containing one or more S, N, O, P, or Si heteroatoms, and the foregoing C_{1-40} hydrocarbyl or C_{3-40} heteroatom containing groups containing a crosslinkable group, with the proviso that in at least one occurrence, R^1 comprises crosslinkable group.
- 3. A compound according to claim-1 wherein R¹ in at least one occurrence contains a double bond, a triple bond, a precursor capable of in situ formation of a double bond, or a heterocyclic, addition polymerizable group.

5

10

15

20

4. A compound according to claim 1 wherein R¹ in at least one occurrence is selected from the group consisting of:

$$-(R^{5})_{m}-CR^{4}=CR^{4}_{2}, -(R^{5})_{m}-C\equiv CR^{4}, -(R^{5})_{m}-O(R^{5})_{m} CR^{4}=CR^{4}_{2}, -(R^{5})_{m}-O(R^{5})_{m} C\equiv CR^{4}, \\ -(R^{5})_{m}-C(O)(R^{5})_{m} CR^{4}=CR^{4}_{2}, -(R^{5})_{m}-C(O)(R^{5})_{m} C\equiv CR^{4}, -(R^{5})_{m}-OC(O)(R^{5})_{m} CR^{4}=CR^{4}_{2}, \\ -(R^{5})_{m}-OC(O)(R^{5})_{m} C\equiv CR^{4}, -(R^{5})_{m}-COO(R^{5})_{m} CR^{4}=CR^{4}_{2}, -(R^{5})_{m}-COO(R^{5})_{m} C\equiv CR^{4}, \\ -(R^{5})_{m}-O(CO)O(R^{5})_{m} CR^{4}=CR^{4}_{2}, -(R^{5})_{m}-O(CO)O(R^{5})_{m} C\equiv CR^{4},$$

$$NR^4$$
, NR^5 , NR^5 , and NR^4

where

 R^4 is hydrogen, halogen, C_{1-20} hydrocarbyl, C_{1-20} halohydrocarbyl, or C_{1-20} halocarbyl; R^5 is C_{1-20} hydrocarbylene, C_{1-20} halohydrocarbylene, or C_{1-20} halocarbylene; and m is 0 or 1.

- 5. A compound according to claim 1 wherein R¹ is selected from the group consisting of: vinyl, C₁₋₄ alkylacrylate, vinylphenyl, vinylphenyloxy, maleimido, vinylbenzyl, vinylbenzyloxy, oxetanyl, 2-propynyl, trifluoroethenyl, 1-benzo-3,4-cyclobutane, and methyl-1-benzo-3,4-cyclobutane.
- 6. A compound according to claim 1 wherein R² independently each occurrence is hydrogen, C₁₋₂₀ hydrocarbyl, C₁₋₂₀ halohydrocarbyl, C₁₋₂₀ halocarbyl, C₁₋₂₀ hydrocarbyloxy, C₁₋₂₀ hydrocarbyloxy, C₁₋₂₀ hydrocarbyloxy, C₁₋₂₀ hydrocarbyloxy, or cyano.
 - 7. A compound according to claim 6 wherein R² each occurrence is hydrogen.
- 25 8. A compound according to claim 1 wherein Ar¹, Ar², Ar³ and Ar⁴ are phenyl or phenylene, X¹ and X² are O or S, and a and b are 0 or 1.
 - 9. An oligomer or polymer having one or more repeating groups of the formula:

5

10

15

20

25

$$(X^{1})_{a}$$
 Ar^{2}
 R^{*}
 R^{*}
 R^{*}
 Ar^{4}
 $(X^{2})_{b}$
(Ia)

wherein R^* is independently in each occurrence i) a C_{1-40} hydrocarbyl group, iii) a C_{1-40} hydrocarbyl group wherein one or more carbons are substituted by one or more heteroatoms selected from S, N, O, P, B or Si atoms, or iii) a halogenated derivative of i) or ii), with the proviso that in at least one occurrence, R^* is a divalent linking group formed by crosslinking of a crosslinkable group selected from i), ii) or iii) through which the repeating groups are joined;

 R^2 is independently in each occurrence hydrogen, halogen, C_{1-20} hydrocarbyl, C_{1-20} hydrocarbyloxy, C_{1-20} thioether, C_{1-20} hydrocarbylcarbonyloxy, di(C_{1-20} hydrocarbyl)amino, or cyano;

 Ar^1 , Ar^2 , Ar^3 and Ar^4 are independently in each occurrence $C_{6\cdot 20}$ aromatic groups, optionally containing one or more S, N, O, P, B or Si heteroatoms, halo-, $C_{1\cdot 20}$ hydrocarbyl-, $di(C_{1\cdot 20}$ hydrocarbyl)amino-, $C_{1\cdot 20}$ hydrocarbyloxy-, $tri(C_{1\cdot 10}$ hydrocarbyl)silyl-, or $tri(C_{1\cdot 10}$ hydrocarbyl)siloxy- substituted derivatives thereof, or divalent derivatives of the foregoing;

a and b independently in each occurrence are 0 or 1; and

 X^1 and X^2 independently in each occurrence are a covalent bond, O, S, SO₂, CH₂, C(R³)₂ or NR³, wherein R³ is selected from the group consisting of C₁₋₂₂ alkyl, C₁₋₂₂ cycloalkyl, C₆₋₂₄ aryl, and C₇₋₂₄ aralkyl.

- 10. A composition comprising an oligomer or polymer according to claim 9.
- 11. A process for preparing oligomers or polymers comprising heating a composition according to claim 1 under reaction conditions sufficient to form an oligomer or polymer having one or more groups according to claim 9.
 - 12. A composition according to claim 9 in the form of a film.
- 13. An electronic device comprising one or more layers of polymer films, at least one of which comprises a film according to claim 12.
- 30 14. An electronic device according to claim 13 which is an electroluminescent device.